



02-15-00

A



In re U.S. Continuation application of: UNIVERSITY OF LEICESTER

Inventor: Wilhelm SCHWAEBLE

For: C1q AND COLLECTIN RECEPTOR



**\*\*THIS APPLICATION CLAIMS PRIORITY FROM PCT/GB98/02430 (FILED AUGUST 12, 1998) AND BRITISH APPLN. NO. 9716998.1 (FILED AUGUST 12, 1997)\*\***

Our Docket No. 3523 P 004

ENCLOSED:

POSTCARD

REQUEST FOR FILING A CONTINUATION OF AN INTERNATIONAL APPLICATION

PRELIMINARY AMENDMENT

INTERNATIONAL APPLICATION AS PUBLISHED, WITH SEARCH REPORT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

TO FOLLOW:

EXECUTED DECLARATION/POWER OF ATTORNEY

CHECK FOR PAYMENT OF FILING FEE

VERIFIED STATEMENT FOR SMALL ENTITY

I HEREBY CERTIFY THAT THIS PAPER AND THE ABOVE DOCUMENTS ARE BEING DEPOSITED WITH THE U.S. POSTAL SERVICE AS EXPRESS MAIL IN AN ENVELOPE ADDRESSED TO BOX PATENT APPLICATION/NO FEE, ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, DC 20231, ON February 14, 2000 UNDER EXPRESS MAIL NO. EE486969926US.

*Husan M. Frank*

**PLEASE ADDRESS ALL FUTURE COMMUNICATIONS TO:**

**MONIQUE A. MORNEAULT, ESQ.  
WALLENSTEIN & WAGNER  
311 South Wacker Drive - 5300  
Chicago, IL 60606  
(312) 554-3300**

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In Re U.S. Patent Application of: )  
Wilhelm SCHWAEBLE )  
 )  
From: PCT/GB98/02430 (filed 8/12/98) )  
and British Appln. No. 9716998.1 (filed 8/12/97) )  
 )  
For: C1q AND COLLECTIN RECEPTOR )  
 )

**REQUEST FOR FILING A CONTINUATION OF AN INTERNATIONAL APPLICATION**

ASSISTANT COMMISSIONER FOR PATENTS  
Washington, D.C. 20231

ATTN: BOX PATENT APPLICATION/NO FEE

Sir:

This is a request for filing a continuation application under 37 C.F.R. 1.53(b), of pending prior International Application No. PCT/GB98/02430, filed August 12, 1998 (claiming priority from British Appln. No. 9716998.1, filed August 12, 1997), titled C1q AND COLLECTIN RECEPTOR, which designated the United States.

Enclosed are the specification, claims and Abstract, along with five sheets of formal drawings (Figures 1-5).

Amend the specification by inserting before the first line the sentence: "This Application is a Continuation of International Application No. PCT/GB98/02430, filed August 12, 1998 (claiming priority from British Appln. No. 9716998.1, filed August 12, 1997), now pending (which is hereby incorporated by reference)."

⇒ A Preliminary Amendment is enclosed.

Attorney Docket No. 3523 P 004  
Title: C1q AND COLLECTIN RECEPTOR

Page 2

We also enclose for completeness:

- ⇒ International application as published, with Search Report; and
- ⇒ International Preliminary Examination Report.

Respectfully submitted,

Date: February 14, 2000

By: Monique A. Morneault  
Monique A. Morneault Reg. No. 37,893  
WALLENSTEIN & WAGNER, LTD.  
311 South Wacker Drive - 5300  
Chicago, IL 60606  
1-312-554-3300

smf 91646

Express Mail Label No. EE486969926US

Date of Deposit: FEBRUARY 14, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to:  
ASSISTANT COMMISSIONER FOR PATENTS, Washington, D.C. 20231  
(ATTN: BOX PATENT APPLICATIONS/NO FEE).

Lusan M. Frankh

In Re U.S. Patent Application of:  
Wilhelm SCHWAEBLE

From: PCT/GB98/02430 (filed 8/12/98)  
and British Appln. No. 9716998.1 (filed 8/12/97)

For: C1q AND COLLECTIN RECEPTOR

## PRELIMINARY AMENDMENT

ASSISTANT COMMISSIONER FOR PATENTS  
Washington, D.C. 20231

ATTN: BOX PATENT APPLICATION/NO FEE

Please amend the above-identified application as follows:

IN THE SPECIFICATION:

Amend the specification by inserting before the first line the sentence: "This Application is a Continuation of International Application No. PCT/GB98/02430, filed August 12, 1998 (claiming priority from British Appln. No. 9716998.1, filed August 12, 1997), now pending (which is hereby incorporated by reference)."

IN THE CLAIMS:

Claim 1, line 1, after "The" delete "use of" and insert - - A molecule comprising - -; after "domain" insert - - - useful - -.

Claim 2, line 1, after "The" delete "use of a cClqR binding domain" and insert - - molecule-; after "the" insert - - cClqR - -.

Claim 3, line 1, after "The" delete "use of a cC1qR binding domain" and insert - - molecule - -; lines 1-2, delete "either one of claims 1 or 2" and insert - - claim 1 - -.

Claim 4, delete "The use of" and insert - - A molecule comprising - -; after "domain" insert - - useful - -.

Claim 5, after "The" delete "use of an inhibitor of the c1qR binding domain" and insert - - molecule - -; after "claim 4" insert - - wherein - -; line 2, delete "comprising" and insert - - comprises - -.

Claim 6, line 1, after "The" delete "use of an inhibitor of the cC1qR binding domain according to either one of claims 4 or 5" and insert - - molecule according to claim 4, wherein the inhibitor of the cC1qR binding domain is useful - -.

Claim 7, line 1, after "The" delete "use of an inhibitor of the cC1qR domain" and insert - - molecule - -; after "claim 6" insert - - wherein the inhibitor of the cC1qR binding domain is useful - -.

Claim 8, line 1, after "The" delete "use of an inhibitor of the cC1qR binding domain according to either one of claims 4 or 5" and insert - - molecule according to claim 4, wherein the inhibitor of the cC1qR binding domain is useful - -.

Claim 9, lines 1-2, delete "The use of a cC1qR domain or inhibitor thereof according to any one of the preceding claims" and insert - - A molecule comprising a cC1qR binding domain, - -.

Claim 10, line 2, after "inhibitor thereof," delete "according to any one of the preceding claims".

No new matter is being added through these amendments. Applicant respectfully requested entry of the above amendments.

Respectfully submitted,

Date: February 14, 2000

By: Monique A. Morneault  
Monique A. Morneault Reg. No. 37,893  
WALLENSTEIN & WAGNER, LTD.  
311 South Wacker Drive - 5300  
Chicago, IL 60606  
1-312-554-3300

smf 91893

Express Mail Label No. EE486969926US

Date of Deposit: FEBRUARY 14, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. § 1.10 on the date indicated above and is addressed to:  
ASSISTANT COMMISSIONER FOR PATENTS, Washington, D.C. 20231  
(ATTN: BOX PATENT APPLICATIONS/NO FEE).

Luis A. M. Frank

### C1q and Collectin Receptor

The present invention concerns novel uses of the C1q and collectin Receptor (cC1qR) binding domain and inhibitors thereof.

The C1q binding domain within cC1qR (cC1q receptor) has previously been identified (Stuart, G.R. *et al.*, 1996, FEBS Letters, 397: 245-249 and references therein). The cC1q receptor is also referred to as the C1q receptor, which also binds collectins. The cC1qR homologue Calreticulin (CaR/CRT) has also been identified and shown to have very high sequence homology, and as such reference to cC1qR is considered to also be reference to CRT and *vice versa* unless otherwise stated.

Certain functionality has been attributed to cC1qR upon its activation by the binding of C1q, namely immunological responses such as phagocytosis, enhanced cytokine and antibody production and antibody-dependent cell cytotoxicity. cC1qR is also known to bind the collectin proteins SP-A, MBL, CL43 and conglutinin. However, the exact nature of cC1qR has not yet been determined nor its structure identified. Sequence analysis does not identify it as being part of a known class of cell-surface receptors.

The present inventors have now found that the C1q binding domain of cC1qR is in fact a CUB (Complement Ubiquitin) domain, and as such certain previously unknown functionality can be attributed to cC1qR and inhibitors of same. CUB domains are well known (see for example Day, A.J. *et al.*, 1993, Behring Inst. Mitt., 93: 31-40; Thiel, S. *et al.*, 1997, Nature, 386: 506-510; Arlaud, G.J. *et al.*, 1993, Behring Inst. Mitt., 93: 189-195).

Thus according to the present invention there is provided the use of a cC1qR binding domain as a CUB domain. Thus the present invention may provide the use of a cC1qR binding domain in the manufacture of a medicament to effect CUB domain functionality. Since the cC1qR binding domain is both a collectin and C1qR binding domain, the term "cC1qR binding domain" also encompasses C1q binding domains.

Sequence analysis of cC1qR shows little primary homology with known CUB domains, but the cC1qR binding domain does in fact have 6 to 7 consensus residues (out of a total of about 100) with CUB and this provides the cC1qR binding domain (and thus cC1qR) with CUB domain functionality.

The C1q binding domain may form part of an existing molecule, for example cC1qR or it may form part or the whole of a novel molecule, for example a molecule comprising a recombinant cC1qR binding domain. The cC1qR binding domain may bind a site comprising five collagen repeats (Gly-X-Y triplets) (Malhotra, R. *et al.*, Biochem. J., 293: 15-19).

Also provided according to the present invention is the use of an inhibitor of the cC1qR binding domain to inhibit CUB functionality. Such an inhibitor may of course be any molecule or other chemical agent which is capable of inhibiting the activation of the cC1q receptor. Examples of such inhibitors include recombinant cC1qR binding domains which competitively inhibit the binding of C1q to cC1qR and thereby inhibit the activation of the cC1q receptor.

Such competitive inhibitors (a) inhibit C1q and mannose binding lectin mediated activation of complement and (b) bind to the part of the collagenous region of C1q where covalent binding to ss-amyloid occurs in Alzheimer's disease.

- 3 -

Also provided according to the present invention is a method of manufacture of a medicament to effect CUB domain functionality, characterised in the use of a cC1qR binding domain. Also provided according to the present invention is a method of manufacture of a medicament for inhibiting CUB domain functionality, characterised in the use of an inhibitor of the cC1qR binding domain.

The identification of the cC1qR binding domain as a CUB domain provides a wide range of previously unidentified functionality for the cC1q receptor and inhibitors thereof. Inhibition of the cC1qR binding domain prevents (i.e. inhibits) complement activation *via* the classical and lectin pathways and provides therapeutic potential in all such diseases in which complement activation is involved in the initiation and maintenance of inflammation, for example myocardial infarction, brain ischemia (stroke), gut ischemia, rheumatoid arthritis, systemic lupus erythematosus, burns and immune complex nephritis. The cC1qR binding domain may also be used to inhibit the binding of  $\beta$ -Amyloid to C1q, thereby inhibiting the formation of amyloid plaques in Alzheimers disease (Velazquez, P. *et al.*, 1997, Nat. Med., 3(1): 77-79). Additional CUB functionality includes the ability to bind carbohydrate domains of molecules, for example of collagens, and to cause opsonisation.

Thus the present invention also provides the use of an inhibitor of the cC1qR binding domain according to the present invention in the manufacture of a medicament for the treatment of complement activation involved in the initiation and maintenance of inflammation, for example for the treatment of myocardial infarction, brain ischemia (stroke), gut ischemia, rheumatoid arthritis, systemic lupus erythematosus, burns, or immune complex nephritis. Also provided is the use of an inhibitor of the C1qR binding domain according to the present invention in the manufacture of a medicament for the treatment of amyloid plaques in Alzheimers disease.



The present invention also provides the advantage of achieving the desired effecting or inhibiting of CUB domain functionality through the use of a low molecular weight molecule.

Various cC1qR binding domains have been identified by the present inventors, namely those of humans (SEQ ID NO: 1), mice (SEQ ID NO: 2) (*Mus musculus* C57 black) and rats (SEQ ID NO: 3) (*Rattus norvegicus* whistar). Thus the cC1qR binding domain may have the sequence of any one of SEQ ID NOs: 1-3. Obviously, the sequence may be partially modified to retain CUB domain functionality yet have a sequence which is different from the one from which it was derived, i.e. one of SEQ ID NOs: 1-3, and the present invention encompasses the use of such partially modified domains. Partial modification may, for example, be by way of addition, deletion or substitution of amino acid residues. Substitutions may be conserved substitutions. Hence the partially modified molecule may be a homologue of the molecules from which it was derived. It may, for example, have at least 40% homology with the molecule from which it was derived. It may for example have at least 50, 60, 70, 80, 90 or 95% homology with the molecule from which it was derived. An example of a homologue is an allelic mutant.

Also provided according to the present invention is a method of treatment of the human or animal body comprising the use of a cC1qR binding domain or an inhibitor thereof according to the present invention. The method may be a method for effecting (in the case of a cC1qR binding domain) or the inhibiting (in the case of a cC1qR binding domain inhibitor) CUB domain functionality.

The invention will be further apparent from the following description, with reference to the several figures of the accompanying drawings, which show, by way of example only, forms of cC1qR binding domain. Of the figures:

- 5 -

Figure 1 shows the amino acid sequence of CRT (SEQ ID NO: 7). Amino acid sequences were deduced from the nucleotide sequence of human CRT (McCauliffe, D.P. *et al.*, 1990, J. Clin. Invest., 85: 1379-1391.). The signal sequence residues are shown in lower case. The N- (*italics*), P- and C- (*italics*) domains are indicated. The CUB-domain is underlined. Domain constructs were expressed as thioredoxin fusion products;

Figure 2 shows binding of C1q to Calreticulin domains. Solid-phase bound domains, with appropriate controls, were incubated with radioiodinated C1q. Binding levels of four separate experiments, at saturation, are shown, calculated as % (bound/loaded). These percentages were then standardised against the results for cC1qR. Y-axis shows % relative binding. X-axis shows immobilised proteins (left to right): C1qR, N-domain, S-domain, P-domain, C-domain and BSA;

Figure 3 shows binding of CUB-domain to immobilised C1q. Serial dilutions of radiolabelled CUB-domain were bound to immobilised C1q and BSA. After extensive washing, bound radioactivity was measured as described in Experimental section (below). Y-axis shows cpl bound. X-axis shows cpm loaded. Upper line (solid, with open squares) is C1q. Lower line (dashed, with open circles) is BSA;

Figure 4 shows the inhibition of CUB-domain-C1q interaction by collectins, C1q and C1q collagen tails. Constant levels of radiolabelled CUB-domain were pre-incubated with serial dilutions of unlabelled C1q, C1q tails and collectin proteins. The incubation mixture was bound to, and eluted from, solid phase C1q. Solid line with open squares is C1q; dashed line with diamonds is SP-A; dashed line with open triangles is MBL; solid line with solid circles is SP-D; dashed line with solid squares is CL43; dashed line with open squares is C1q tails; solid line with open circles is BSA; and

- 6 -

Figure 5 shows the inhibition of the classical pathway of complement activation by addition of the recombinant CUB domain of cC1qR. Haemolytic assays were used to determine the concentration-dependence of the inhibition of the classical activation pathway of complement by the recombinant CUB domain, the N-domain and the P-domain of cC1qR. Y-axis shows % inhibition. X-axis shows  $\mu\text{g/ml}$ . Line with solid circles is CUB-domain. Line with solid squares is P-domain. Line with solid hexagons is N-domain. Line with open circles is BSA.

## EXPERIMENTAL

The following experiments demonstrate the activity of the C1q binding domain of cC1qR and that the C1q binding domain of cC1qR is in fact a CUB and show (see Figure 5) the inhibition of the classical pathway of complement activation by addition of the recombinant CUB domain of cC1qR.

The sequence listing includes: the DNA sequence (SEQ ID NO: 4) and derived amino acid sequence (SEQ ID NO: 1) of human calreticulin; the DNA sequence (SEQ ID NO: 5) and derived amino acid sequence (SEQ ID NO: 2) of mouse calreticulin; the DNA sequence (SEQ ID NO: 6) and derived amino acid sequence (SEQ ID NO: 3) of rat calreticulin; and the amino acid sequence of CRT (SEQ ID NO: 7). Amino acid sequences were deduced from the nucleotide sequence of human CRT (McCauliffe, D.P. *et al.*, 1990, J. Clin. Invest., 85: 1379-1391.). The signal sequence residues of SEQ ID NO: 7 are residues 1-17. The N-domain (shown in *italics*) is residues 18-196; the P- (residues 197-308) and C- (residues 309-417) (*italics*) domains are indicated. The CUB-domain is residues 162-283 (underlined). Domain constructs were expressed as thioredoxin fusion products.

Recombinant CRT domains to be tested for C1q and collectin binding function were produced from a cDNA clone (phCRT-1) isolated from a human umbilical vein endothelial cell cDNA library (Stuart, G.R. *et al.*, 1996, Exp. Lung Res., 22: 467-487; Stuart, G.R. *et al.*, 1996, FEBS Lett., 397: 245-249). These domains, as described below, are based upon structural predictions for the molecule and have previously been used to localise CRT function within the molecule. The amino-terminal N-domain contains the binding regions for PDI (Baksh, S. *et al.*, 1995, J. Biol. Chem., 270(52): 31338-31344), Zn<sup>2+</sup> (Baksh, S. *et al.*, 1995, FEBS Lett., 376(1-2): 53-57) and integrins (Leung-Hagesteijn, C.Y. *et al.*, 1994, J. Cell Sci., 107 (Pt 3): 589-600). The proline-rich central P-domain contains the high affinity Ca<sup>2+</sup> binding site (Baksh, S. and Michalak, M., 1991,

J. Biol. Chem., 266: 21458-21465) and the lectin site (D. Williams, cited in Krause, K.H. and Michalak, M., 1997, Cell, 88(4): 439-443) within two sets of highly conserved repeats. The acidic C-domain contains the ER-retention terminal KDEL signal (McCauliffe, D.P. *et al.*, 1990, J. Clin. Invest., 85: 1379-1391) and the low affinity Ca<sup>2+</sup> binding site (Baksh, S. and Michalak, M., 1991, J. Biol. Chem., 266: 21458-21465). Previous studies have indicated that the C1q binding site lies across the intersection of the N and P-domains (Stuart, G.R. *et al.*, 1996, FEBS Lett., 397: 245-249; Stuart, G.R. *et al.*, 1997, Immunopharmacology, 38:73-80). Within this region we have identified and expressed a 123 amino acid region containing a putative C1r/C1s (also termed CUB) module (Day, A.J. *et al.*, 1993, Behring Inst. Mitt., 93: 31-40) based upon amino acid sequence alignments. We termed this segment the CUB-domain and show here that it contains the C1q and collectin-binding site of cC1qR/CRT.

#### *Purification and Radioiodination of cC1qR, C1q and Collectins*

Native cC1qR was purified from human U937 cells as previously described (Malhotra, R. *et al.*, 1993, Immunology, 78: 341-348). cC1qR and CUB-domain samples were iodinated by the Iodogen method (Fraker, P.J. and Speck, J.C. Jr., 1978, Biochem. Biophys. Res. Commun., 80: 849-857). C1q was purified as previously described (Reid, K.B.M., 1981, Methods in Enzymology, 80: 16-25) and radioiodinated as described by Bolton & Hunter (Bolton, A.E. and Hunter, W.M., 1973, Biochem. J., 133: 529-539) as this method of iodination causes less damage to large, oxidation-sensitive molecules such as C1q than the more frequently utilised Iodogen method (Stuart, G.R. *et al.*, 1996, Exp. Lung Res., 22: 467-487). Radiolabelled proteins were stored at 4 °C. C1q collagen tails were prepared as described by Reid (Reid, K.B.M., 1976, Biochem. J., 155: 5-17). Collectins were purified as previously described (Malhotra, R. *et al.*, 1990, J. Exp. Med., 172: 955-959; Holmskov, U. *et al.*, 1995, Biochem. J., 305: 889-896).

#### *Prokaryotic expression of recombinant Calreticulin domains*

Given that CRT, Clr and Cls all interact with Clq, a sequence comparison was performed to investigate the structural basis for this interaction. A region that may correspond to a CUB module was identified in CRT and was analysed by multiple sequence alignment as described previously (Day, A.J. *et al.*, 1993, Behring Inst. Mitt., 93: 31-40). This region, termed the S region (Cls-like (CUB) domain), spans the intersection of the N and P-domains (residues 160-283). A 1.9kb cDNA clone for CRT (phCRT-1) was isolated from a human umbilical vein endothelial cell library in the eukaryotic expression vector CDM8 (Aruffo, A. and Seed, B., 1987, PNAS USA, 84: 8573-8577). Sequence analysis revealed that phCRT-1 comprised the complete coding sequence for CRT with absolute identity to the previously published human CRT sequence (McCauliffe, D.P. *et al.*, 1990, J. Clin. Invest., 85: 1379-1391).

The Thiobond expression system was used to produce N, P, C and CUB-domains of cClqR/CRT (representing the N-terminal region, the proline-rich central region, the C-terminal region, and a region spanning the intersection of the N and P-domains (as described above) (Figure 1). The individual domains were expressed as thioredoxin fusion proteins in *E. Coli* using the plasmid pTrxfus (Invitrogen BV, Leek, Netherlands) as described previously (Stuart, G.R. *et al.*, 1996, FEBS Lett., 397: 245-249; Stuart, G.R. *et al.*, 1997, Immunopharmacology, 38:73-80).

Samples were assayed for recombinant calreticulin domain expression by SDS-PAGE (Laemmli, U.K., 1970, Nature, 227: 680-685) and by Western blotting with rabbit antisera to: (1) whole cClqR (raised against human cClqR purified from U937 cells (Malhotra, R. *et al.*, 1993, Immunology, 78: 341-348); (2) CRT C-terminal region (raised against a GST fusion protein containing the final 18 residues of recombinant human CRT), and (3) CRT N-terminal region (raised against a GST fusion protein containing residues 7-18 of recombinant human CRT).

- 10 -

*Interaction of immobilised recombinant human cC1qR/CRT domains with radiolabelled C1q*

Binding experiments with the cC1qR/CRT domains were performed throughout in low salt (10mM potassium phosphate, 0.5mM EDTA (pH 7.4)) in order to maximise the ionic interaction with C1q.

Microtitre plates were coated with the N-, P-, C- and CUB-domains and with three controls, cC1qR, BSA and Thioredoxin, (8mg/ml in 35mM NaHCO<sub>3</sub>, 15mM Na<sub>2</sub>CO<sub>3</sub> pH 9.6) for 2hr at 37 °C. Non-specific interactions were blocked by incubation with 10mM potassium phosphate, 0.5mM EDTA pH 7.4 containing BSA (10mg/ml). Any free -SH groups in the samples, due to the presence of the thioredoxin fusion protein, were blocked by a brief washing step using the phosphate buffer containing 2mM iodoacetamide. After washing, serial dilutions of radioiodinated C1q (in 10mM potassium phosphate, 0.5mM EDTA, pH 7.4) were added to the wells and incubated for 2h at 37 °C. Wells were washed three times with phosphate buffer and bound radioactivity eluted with 100ml 4M NaOH and measured.

*Concentration dependent binding of radiolabelled CUB-domain to immobilised C1q*  
*C1q binds to the Fc regions of IgG.*

This property was utilised in order to correctly orient the C1q on microtitre plates. Breakable microtitre plates (Life Sciences International) were coated with rabbit Fc (5mg per well in 35mM NaHCO<sub>3</sub>, 15mM Na<sub>2</sub>CO<sub>3</sub> pH 9.6) . Non-specific sites were blocked as described above, and the wells were incubated with C1q (5 mg per well in 10mM potassium phosphate buffer). Certain wells were also coated with BSA as a negative control. After further washing, serial dilutions of radioiodinated CUB-domain (in 10mM potassium phosphate, 0.5mM EDTA, pH 7.4) were added to the wells and incubated for 2h at 37 °C. Wells were washed three times with the phosphate buffer and bound radioactivity in the individual wells measured.

*Competitive inhibition of the CUB-domain-C1q interaction by fluid phase C1q, collectins and C1q collagen tails*

C1q was immobilised onto Fc-coated microtitre plates as described above. Non-specific binding was blocked by incubation with 10mM potassium phosphate, 0.5mM EDTA (pH 7.4) containing BSA (10mg/ml). Serial dilutions of the collectins (SP-A, MBL, SP-D, CL43), C1q, C1q tails and BSA (maximum quantity = 9mg/well) were prepared in 10mM potassium phosphate, 0.5mM EDTA (pH 7.4). Each dilution (100µl) was then incubated for 1h at 37 °C with a constant level of radiolabelled CUB-domain and loaded onto the plate. Following 2 hours incubation at 37 °C, wells were extensively washed and bound radioactivity measured.

## Results

### *Amino acid sequence alignments*

A region spanning the intersection of the N- and P-domains of CRT has previously been implicated in C1q binding (Stuart, G.R. *et al.*, 1996, FEBS Lett., 397: 245-249; Stuart, G.R. *et al.*, 1997, Immunopharmacology, 38:73-80). Amino acid sequence alignment of a region within CRT (residues 160-283) showed little primary homology to a CUB module, but the region is in fact a CUB module. Two CUB modules, together with an EGF module, form a binding region within C1r2C1s2 for the collagenous tails of C1q. cC1qR competes with C1r2C1s2 for binding to C1q, implying a similarity in C1q binding sites on C1r, C1s and cC1qR (Sobel, A.T. and Bokisch, V.A., in: Membrane receptors of lymphocytes (M. Seligman, FL Preud'homme, FM Kourilsky eds). North Holland Publishing Co., Amsterdam, p151, 1975; Van den Berg, R.H. *et al.*, 1995, Eur. J. Immunol., 25(8): 2206-2210). This segment, the CUB-domain (residues 162-283, see Figure 1), was tested for C1q binding.

### *Interaction of recombinant human Calreticulin domains with C1q*

N-, P-, C-, and CUB-domains of human CRT were expressed as thioredoxin fusion proteins. Correct expression was verified by SDS-PAGE and Western blotting. Figure



2 summarises the results of four separate solid phase direct binding experiments. Significant binding to radioiodinated C1q was observed for cC1qR, the CUB-domain, the P-domain, and, to a lesser extent, the N-domain. The C-domain showed no binding. Serial dilutions of radioiodinated CUB-domain were incubated with immobilised C1q and BSA (Figure 3). Concentration-dependent, saturable binding was observed to C1q but not to BSA.

*Competitive inhibition of C1q-CUB-domain interaction by C1q tails and Collectins.*

C1q was immobilised on microtitre plates by interaction with solid phase Fc. Figure 4 shows the results of competitive inhibition of the CUB-domain-C1q interaction. As expected, native fluid-phase C1q demonstrates concentration-dependent inhibition. C1q tails also cause inhibition, indicating that the interaction of the CUB-domain with C1q is via the collagenous C1q tails. Inhibition studies with the collectin proteins demonstrated that SP-A, MBL and CL43 interact with the CUB-domain, via the same, or an overlapping binding site as C1q. SP-D and BSA did not inhibit the CUB-domain-C1q interaction.

*Competitive inhibition of C1q mediated complement activation, i.e. classical pathway activation.*

The ability of cC1qR and fragments thereof to bind to C1q and interfere with binding of the C1q associated serine proteases C1r2C1s2 was determined by haemolysis assays. Sheep erythrocytes (E) and rabbit anti-sheep erythrocytes (A) were used to prepare EAC1q cells as described by Borsos and Rapp (Borsos, T. and Rapp, H.J., 1967, J. Immunol., 99: 263-268) and de Bracco and Stroud (De Bracco, M.M. and Stroud, R.M., 1971, J. Clin. Invest. 50: 838-848). EA and EAC1q cells ( $100\text{ml} \times 10^8$  cells/ml) were incubated at 37 °C for 1h with serial dilutions of C1q deficient serum, from a patient with homozygous C1q deficiency, in order to establish the minimum serum concentration required to cause complete cell lysis of EAC1q cells. EAC1q cells (100ml) were then incubated for 1h at 37 °C with increasing concentrations of the recombinant N-domain,

- 13 -

P-domain and the CUB-domain of human cC1qR and BSA as a control and after incubation, the extent of lysis established by the addition of cold DGVB<sup>++</sup> (1ml) (Mayer, 1961). After centrifugation (3000g, 10 min) the OD<sup>412</sup> of the supernatant was measured. Controls of 100% lysis comprise 100ml cells and 1.1 ml water. The results of this assay are shown in Figure 5. Hereby, the CUB domain of cC1qR was shown to effectively inhibit C1q mediated complement activation. At higher concentrations, the P-domain and the N-domain (which contain parts of the CUB-domain) show inhibition at a lesser extent.

CLAIMS

1. The use of a cC1qR binding domain in the manufacture of a medicament to effect CUB domain functionality.
2. The use of a cC1qR binding domain according to claim 1, the binding domain forming part of a C1q receptor.
3. The use of a cC1qR binding domain according to either one of claims 1 or 2, the cC1qR binding domain being recombinant.
4. The use of an inhibitor of the cC1qR binding domain in the manufacture of a medicament to inhibit CUB functionality.
5. The use of an inhibitor of the cC1qR binding domain according to claim 4, the inhibitor comprising a cC1qR binding domain.
6. The use of an inhibitor of the cC1qR binding domain according to either one of claims 4 or 5 in the manufacture of a medicament for the treatment of complement activation involved in the initiation and maintenance of inflammation.
7. The use of an inhibitor of the cC1qR binding domain according to claim 6 in the manufacture of a medicament for the treatment of myocardial infarction, brain ischemia (stroke), gut ischemia, rheumatoid arthritis, systemic lupus erythematosus, burns, or immune complex nephritis.

- 15 -

8. The use of an inhibitor of the cC1qR binding domain according to either one of claims 4 or 5 in the manufacture of a medicament for the treatment of amyloid plaques in Alzheimers disease.

9. The use of a cC1qR binding domain or inhibitor thereof according to any one of the preceding claims, the cC1qR binding domain having the sequence of any one of SEQ ID NOs: 1-3 or a partially modified form thereof.

10. A method of treatment of the human or animal body comprising the use of a cC1qR binding domain or an inhibitor thereof according to any one of the preceding claims.


1 / 5

## FIGURE 1

1 mllsvplllg llglavaEPA VYFKEQFLDG DGWTPRWIES KHKSDFGKFV  
51 LSSGKFYGD EKD KGLQTSQ DARFYALSAS FEPFSNKGQT LVVQFTVKHE  
101 QNIDCGGGYV KLFPNSLDQT DMHGDSEYNI MFGPDICGPG TKKVHVIFNY  
151 KGKNVLINKD IRCKDDEFTH LYTLIVRPDN TYEVKIDNSO VESGSLEDDW  
201 DFLPPKKIKD PDASKPEDWD ERAKIDDPTD SKPEDWDKPE HIPDPDAKKP  
251 EDWDEEMDGE WEPPVIONPE YKGEWKPROI DNPDYKGTWI HPEIDNPEYS  
301 PDPSIYAYDN FGVGLDLWQ VKSGTIFDNF LITNDEAYAE EFGNETWGV  
351 KAAEQMKDK QDEEQRLKEE EEDKKRKEEE EAEDKEDDED KDEDEEDED  
401 KEEDEEDVP GQAKDEL


2/5

FIGURE 2




3 / 5

FIGURE 3



4 / 5


FIGURE 4





5/5

FIGURE 5



## SEQUENCE LISTING

<110> SCHWAEBLE, Wilhelm  
University of Leicester, The

<120> Clq and Collectin Receptor

<130> M97/0287/PCT

<140>

<141>

<160> 7

<170> PatentIn Ver. 2.0

<210> 1

<211> 122

<212> PRT

<213> Homo sapiens

<400> 1

Arg Cys Lys Asp Asp Glu Phe Thr His Leu Tyr Thr Leu Ile Val Arg  
1 5 10 15

Pro Asp Asn Thr Tyr Glu Val Lys Ile Asp Asn Ser Gln Val Glu Ser  
20 25 30

Gly Ser Leu Glu Asp Asp Trp Asp Phe Leu Pro Pro Lys Lys Ile Lys  
35 40 45

Asp Pro Asp Ala Ser Lys Pro Glu Asp Trp Asp Glu Arg Ala Lys Ile  
50 55 60

Asp Asp Pro Thr Asp Ser Lys Pro Glu Asp Trp Asp Lys Pro Glu His  
65 70 75 80

Ile Pro Asp Pro Asp Ala Lys Lys Pro Glu Asp Trp Asp Glu Glu Met  
85 90 95

Asp Gly Glu Trp Glu Pro Pro Val Ile Gln Asn Pro Glu Tyr Lys Gly  
100 105 110

Glu Trp Lys Pro Arg Gln Ile Asp Asn Pro  
115 120

<210> 2

<211> 122

<212> PRT

- 2 -

&lt;213&gt; Mus musculus

&lt;400&gt; 2

Arg Cys Lys Asp Asp Glu Phe Thr His Leu Tyr Thr Leu Ile Val Arg  
 1 5 10 15

Gln Asp Asn Thr Tyr Glu Val Lys Ile Asp Asn Ser Gln Val Glu Ser  
 20 25 30

Gly Ser Leu Glu Asp Asp Gly Asp Phe Leu Pro Pro Lys Lys Ile Lys  
 35 40 45

Asp Pro Asp Ala Ala Lys Pro Glu Asp Trp Asp Glu Arg Ala Lys Ile  
 50 55 60

Asp Asp Pro Thr Asp Ser Lys Pro Glu Asp Trp Asp Lys Pro Glu His  
 65 70 75 80

Ile Pro Asp Pro Asp Ala Lys Lys Pro Glu Asp Trp Asp Glu Glu Met  
 85 90 95

Asp Gly Glu Trp Glu Pro Pro Val Ile Gln Asn Pro Glu Tyr Lys Gly  
 100 105 110

Glu Trp Lys Pro Arg Gln Ile Asp Asn Pro  
 115 120

&lt;210&gt; 3

&lt;211&gt; 122

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 3

Arg Cys Lys Asp Asp Glu Phe Thr His Leu Tyr Thr Leu Ile Val Arg  
 1 5 10 15

Pro Asp Asn Thr Tyr Glu Val Lys Ile Asp Asn Ser Gln Val Glu Ser  
 20 25 30

Gly Ser Leu Glu Asp Asp Trp Asp Phe Leu Pro Pro Lys Lys Ile Lys  
 35 40 45

Asp Pro Asp Ala Ala Lys Pro Glu Asp Trp Asp Glu Arg Ala Lys Ile  
 50 55 60

Asp Asp Pro Thr Asp Ser Lys Pro Glu Asp Trp Asp Lys Pro Glu His  
 65 70 75 80

Ile Pro Asp Pro Asp Ala Lys Lys Pro Glu Asp Trp Asp Glu Glu Met  
 85 90 95

- 3 -

Asp Gly Glu Trp Glu Pro Pro Val Ile Gln Asn Pro Glu Tyr Lys Gly  
 100 105 110

Glu Trp Lys Pro Arg Gln Ile Asp Asn Pro  
 115 120

<210> 4  
 <211> 366  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 cggttgcaagg atgatgagtt tacacacctg tacacactga ttgtgcggcc agacaacacc 60  
 tatgaggtga agattgacaa cagccaggtg gagtcaggct ccttgggaaga cgattgggac 120  
 ttcttgccac ccaagaagat aaaggatcct gatgcttcaa aaccggaaga ctgggatgag 180  
 cgggcccaaga tcgatgatcc cacagactcc aagcctgagg actgggacaa gcccagagcat 240  
 atccctgacc ctgatgctaa gaagcccgag gactgggatg aagagatgga cggagagtgg 300  
 gaacccccag tgattcagaa ccctgagtac aagggtgagt ggaagccccg gcagatcgac 360  
 aaccca 366

<210> 5  
 <211> 366  
 <212> DNA  
 <213> Mus musculus

<400> 5  
 cgggtgtaagg atgatgaatt cacacaccta tacacactga ttgtgcggca agacaacacc 60  
 tatgaggtga aaattgacaa cagccaggtg gagtcaggct ccttggagga tgatggggac 120  
 tttctgccac ccaagaagat aaaggaccct gatgctgcca agccggaaga ctgggatgaa 180  
 cgagccaaga tcgatgaccc cacagattcc aagcctgagg actgggacaa gccagagcac 240  
 atccctgacc ctgatgctaa gaagcctgag gactgggatg aagagatgga tggagagtgg 300  
 gaaccaccag tgattcaaaa tcctgaatac aagggcgagt ggaaaccacg tcaaattgac 360  
 aaccca 366

<210> 6  
 <211> 366

- 4 -

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 6

cgggtgtaagg atgatgaatt cacacatcta tacacgctga ttgtgcggcc agacaacacc 60  
 tacgaggtga aaattgacaa cagccaggtg gagtcgggct ccttggagga tgattgggac 120  
 tttctgccgc ccaagaagat taaggatcct gacgctgcca agccagaaga ctgggatgaa 180  
 cgagccaaga ttgatgaccc cacagattcc aagcctgagg actgggacaa gccagagcac 240  
 atccctgacc ctgatgctaa gaagcctgag gactgggacg aagagatgga tggagagtgg 300  
 gaaccaccag tgattcaaaa tcctgaatac aagggcgaat ggaagccacg tcaaattgac 360  
 aaccca 366

&lt;210&gt; 7

&lt;211&gt; 417

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

Met Leu Leu Ser Val Pro Leu Leu Leu Gly Leu Leu Gly Leu Ala Val  
 1 5 10 15  
 Ala Glu Pro Ala Val Tyr Phe Lys Glu Gln Phe Leu Asp Gly Asp Gly  
 20 25 30  
 Trp Thr Pro Arg Trp Ile Glu Ser Lys His Lys Ser Asp Phe Gly Lys  
 35 40 45  
 Phe Val Leu Ser Ser Gly Lys Phe Tyr Gly Asp Glu Glu Lys Asp Lys  
 50 55 60  
 Gly Leu Gln Thr Ser Gln Asp Ala Arg Phe Tyr Ala Leu Ser Ala Ser  
 65 70 75 80  
 Phe Glu Pro Phe Ser Asn Lys Gly Gln Thr Leu Val Val Gln Phe Thr  
 85 90 95  
 Val Lys His Glu Gln Asn Ile Asp Cys Gly Gly Gly Tyr Val Lys Leu  
 100 105 110  
 Phe Pro Asn Ser Leu Asp Gln Thr Asp Met His Gly Asp Ser Glu Tyr  
 115 120 125  
 Asn Ile Met Phe Gly Pro Asp Ile Cys Gly Pro Gly Thr Lys Lys Val  
 130 135 140

His Val Ile Phe Asn Tyr Lys Gly Lys Asn Val Leu Ile Asn Lys Asp  
 145 150 155 160  
 Ile Arg Cys Lys Asp Asp Glu Phe Thr His Leu Tyr Thr Leu Ile Val  
 165 170 175  
 Arg Pro Asp Asn Thr Tyr Glu Val Lys Ile Asp Asn Ser Gln Val Glu  
 180 185 190  
 Ser Gly Ser Leu Glu Asp Asp Trp Asp Phe Leu Pro Pro Lys Lys Ile  
 195 200 205  
 Lys Asp Pro Asp Ala Ser Lys Pro Glu Asp Trp Asp Glu Arg Ala Lys  
 210 215 220  
 Ile Asp Asp Pro Thr Asp Ser Lys Pro Glu Asp Trp Asp Lys Pro Glu  
 225 230 235 240  
 His Ile Pro Asp Pro Asp Ala Lys Lys Pro Glu Asp Trp Asp Glu Glu  
 245 250 255  
 Met Asp Gly Glu Trp Glu Pro Pro Val Ile Gln Asn Pro Glu Tyr Lys  
 260 265 270  
 Gly Glu Trp Lys Pro Arg Gln Ile Asp Asn Pro Asp Tyr Lys Gly Thr  
 275 280 285  
 Trp Ile His Pro Glu Ile Asp Asn Pro Glu Tyr Ser Pro Asp Pro Ser  
 290 295 300  
 Ile Tyr Ala Tyr Asp Asn Phe Gly Val Leu Gly Leu Asp Leu Trp Gln  
 305 310 315 320  
 Val Lys Ser Gly Thr Ile Phe Asp Asn Phe Leu Ile Thr Asn Asp Glu  
 325 330 335  
 Ala Tyr Ala Glu Glu Phe Gly Asn Glu Thr Trp Gly Val Thr Lys Ala  
 340 345 350  
 Ala Glu Lys Gln Met Lys Asp Lys Gln Asp Glu Glu Gln Arg Leu Lys  
 355 360 365  
 Glu Glu Glu Glu Asp Lys Lys Arg Lys Glu Glu Glu Glu Ala Glu Asp  
 370 375 380  
 Lys Glu Asp Asp Glu Asp Lys Asp Glu Asp Glu Glu Asp Glu Glu Asp  
 385 390 395 400  
 Lys Glu Glu Asp Glu Glu Glu Asp Val Pro Gly Gln Ala Lys Asp Glu  
 405 410 415  
 Leu